



Investigate The Effects Of Folic Acid Supplementation On Homocysteine Levels In Women With Preeclampsia

Elnaz taklifi¹, Alireza Nourazarian^{2*}, Manizheh Sayyah-Melli³, Abdolrasoul Safaiyan⁴

¹ Department of Biology, College of Basic Sciences, Islamic Azad University of Ahar Branch, Ahar, Iran.

² Department of biochemistry and clinical laboratories, Faculty of medicine, Tabriz University of medical sciences, Tabriz, Iran.

³ Women's Reproductive Health Research Center, Tabriz University of Medical Sciences, Tabriz, Iran

⁴ Department of statistics and Epidemiology, Faculty of Health, Tabriz University Medical Science, Tabriz, Iran.

*corresponding author

ABSTRACT

Pre-eclampsia is a syndrome which is characterized by the increase in blood pressure above 140/90, oedema and proteinuria, and is one of the leading causes of maternal death and of complications in pregnancy that can occur in the second half of pregnancy. Approximately 20 % of pregnant women in developing countries and 5 % of pregnant women worldwide are affected, and it is the second leading cause of maternal death in developing countries, including Iran. Homocysteine (2-amino-4-mercapto-butyric acid), is a sulphur-containing amino acid without special codon and is created by an essential amino acid methionine demethylation from the diet. Homocysteine has sulphydryl group, which makes it more susceptible to other thiol oxidation and disulphide formation. Folic acid is water-soluble vitamin, and despite the need to get pregnancy to avoid disruption process of the nervous system of the fetus, NTD for antioxidant effects reduces oxidative stress, and the concentration and oxidation of homocysteine, therefore, diet and supplements during pregnancy is important. This case-control study was conducted on healthy pregnant women visiting Taleghani and Al Zahra hospitals in Tabriz, Iran, as well as on women with pre-eclampsia admitted to these two hospitals. Fifty-eight samples were selected as controls and 58 subjects were selected as cases. The blood samples were prepared with their consents. Parameter measured included homocysteine concentration in blood. In addition, a questionnaire was completed to collect information on age, gestational stage, height, weight, weight before pregnancy and folic acid (B9) intake. The results in this study are based on the inhibition of confounding variables and indicate significant differences in homocysteine status. The people, who received folic acid supplements during pregnancy were 20, and finally 60% of them were diagnosed with gestational hypertension, have been hospitalized, and 40% of them were in the control group. Elevated homocysteine levels can be considered as a factor in the etiology of preeclampsia. The type of diet and dietary antioxidants, including folic acid, in reducing homocysteine-induced oxidative stress, and improve preeclamptic women is important.

Key words: homocysteine, folic acid, pre-eclampsia

INTRODUCTION

Pre-eclampsia is a syndrome which is characterized by the increase in blood pressure above 140/90, oedema and proteinuria, and is one of the leading causes of maternal death and of complications in pregnancy that can occur in the second half of pregnancy (1) Approximately 20 % of pregnant women in developing countries and 5 % of pregnant women worldwide are affected (2), and it is the second leading cause of maternal death in developing countries, including Iran (3) (4) (5) (6) (7). Complications of preeclampsia can be a significant increase in blood pressure, although in the normal range, swelling on the face, hands, and feet in the morning is increased in weight (>450 g/week in the last trimester of pregnancy), and protein in the urine, etc. In severe cases puffiness, blurred vision, headaches, irritability, and abdominal pain are also added; and eclampsia symptoms worsened, stretch muscles, convulsions, and coma can occur (8, 9) Homocysteine (2-amino-4-mercapto-butyric acid), is a sulphur-containing amino acid without special codon and is created by an essential amino acid methionine demethylation from the diet. Homocysteine has sulphhydryl group, which makes it more susceptible to other thiol oxidation and disulphide formation (10) (11) (12) (13) (14). Homocysteine is affected by the amount of protein and increased vitamins, and during exposure to oxidative stress in the body, the use of sulfur-containing amino acid, such as, cysteine and methionine increases because they are converted to homocysteine by trans-methylation and trans sulfuration pathways, i.e., one of the risk factors for cardiovascular disease (15), oxidized homocysteine produces free radicals, such as, superoxide anion (16) (17) (18). ROS (reactive oxygen species) in biological systems are constantly produced by normal metabolism in the body and are necessary for keeping biological balance through various functions. However, overproduction of these molecules and radicals cause damage to the structure, and function of the cell membrane, polyunsaturated fatty acids, thiol groups in proteins, and nucleic acids in DNA. (19) (20) Generally, oxygen radicals are produced continuously in all living organisms and the devastating effects cause cell damage and death; and their production increases in

oxidative conditions. (21) Production of ROS such as anion superoxide ($\text{O}_2^{\bullet -}$), hydroxyl radical ($\text{HO}\cdot$), alkyl peroxide Radical ($\text{ROO}\cdot$) and alcoxyl radicals ($\text{RO}\cdot$) can be enzymatic and non-enzymatic. Mitochondria are the most important cell organelles for the production of radicals, especially $\text{O}_2^{\bullet -}$ and H_2O_2 in mammals (22) (23) (24) Therefore, determination of free radicals, a marker of oxidative stress, is proportional to the condition of the organism may have an important role in homocysteine metabolism (25) (26) (27) Beside damage from free radical-induced oxidative stress conditions, the type of diet during pregnancy and vitamin supplements, such as, folic acid can act as an antioxidant Barrier (28) Since the inverse association is between tHcy and plasma-folate levels, so people who have low-plasma concentrations of vitamin are at risk of hyperhomocysteinemia (29) (30) (31). Folic acid is water-soluble vitamin, and despite the need to get pregnancy to avoid disruption process of the nervous system of the fetus, NTD for antioxidant effects reduces oxidative stress, and the concentration and oxidation of homocysteine; therefore, diet and supplements during pregnancy is important (32) (2) Generally, after receiving a rich diet in methionine, methionine demethylation pathway began. Thus, methionine transforms to the homocysteine with s-adenosyl homocysteine hydrolase, and methyltransferase enzymes and plasma concentration of homocysteine increases, because the way back to this stage of reaction is dependent on the cycle of tetrahydrofolate. So, the importance of folic acid reduces plasma levels of homocysteine are clear. so that the 5-methyl- tetrahydrofolate of tetrahydrofolate cycle renders. Methyl group to homocysteine by demethylation of methionine and homocysteine is converted to methionine in the presence of methionine synthase (33) (34). and obtained tetrahydrofolate converts into 5-methyl tetrahydrofolate in the presence of 5–10 dimethylene tetrahydrofolate and the cycle continues (35) (36) (37) So, folic acid supplementation during pregnancy is important to prevent hyperhomocysteinemia and its complications, including stillbirth, i.e., very low birth weight, premature birth, increased risk of preeclampsia, and neural tube defects in newborns (38) The RDA for folic

acid is 600 µg/day, and it is recommended to have 400 µg/day of B-9. As a supplement and 200 µg/day of diet containing folate, including green vegetables – citrus and fortified foods, such as, bread, cereal foods fortified with folate (1) (39). The present study aims to investigate the effect of the concentration of homocysteine and folic acid supplementation on preeclampsia.

MATERIALS AND METHODS

Sample Selection

This case-control study was conducted based on conscious informed consent of Medical Ethics Committee on 116 pregnant women admitted to teaching hospitals in Tabriz (Taleghani and Al Zahra hospitals) in 2014. and was based on objective selection criteria for participation in the study: not smoking materials drugs, lack of diseases such as diabetes, heart disease and digestive disease, no history of stroke, asthma or thyroid disorders, and not taking medication. The sample included 58 patients with pre-eclampsia; the pre-eclampsia criteria were blood pressure higher than or equal to 140/90 and proteinuria higher than 300 mg per 24 hours' urine. Other questions were asked, such as age, the gestational stage, parity, height, weight before pregnancy, weight at hospitalization and folic acid intake. After getting permission from Tabriz University of Medical Sciences and introduction letters to the hospitals, the admitted women selected were informed of the study and a participation satisfaction and consent questionnaire was completed by them. To assess the condition of the samples, test for homocysteine (hcy) was performed. Fifty-eight pregnant women in the gestational stage of the second half of pregnancy who were referred to the hospital for routine tests were selected as a control group to identify confounding or interfering parameters.

Blood Samples

Full blood samples were taken over three months from two hospitals affiliated with Tabriz University of Medical Sciences. Samples and centrifuged serums between this time period and the measuring time were frozen at a temperature of -20°.

Homocysteine Measurement

The ELISA method was used to measure homocysteine levels. In this method, the separation of free decreased homocysteine to three thiols and then converted it to SAH. After removal of anti-SAH antibody, monoclonal antibody labelled with peroxidase was added, and peroxidase activity of the antioxidant enzymes glutathione peroxidase and superoxide associated with homocysteine concentration was measured using a spectrophotometer absorbed in the samples .

receiving folic acid supplements

At the time of completing the questionnaire containing personal information To evaluate the effect of folic acid on serum levels of homocysteine and preeclampsia Both the patient and control groups receiving folic acid supplements during pregnancy were examined

Statistical analysis

Statistical analysis was performed with SPSS version18.0 software. Statistically comparisons were between the control group and cases. Differences between mean values were carried out using one-way analysis of variance. Data are represented Mean ± SEM. The differences were considered significant when *P < 0.05.

RESULTS

The following table shows details of the two groups' average concentration parameters as measured by the method above.

Table 1: Average concentration parameter measured, including homocysteine in the two groups of controls and case

	Mean in control	Mean in case	Pvalue
HCY	7.458±3.667	11.12±4.511	0.005
HCY: homocysteine			

As shown in the figure 1, there are significant differences in serum homocysteine concentration among the groups of women with and without pre-eclampsia, so that women with pre-eclampsia (11.12±4.511) show higher homocysteine levels than women

without pre-eclampsia (7.458±3.667) (P<0.05).




Figure 1: Mean of homocysteine concentrations between the groups of women with and without pre-eclampsia

Group Statistics

GROUP		Mean	Std. Deviation
SBP.PR	case	146.7500	13.61782
	Control	113.8889	9.40025
DBP.PR	case	93.3621	9.29006
	Control	74.2593	6.89595

Std: standard Deviation

Table 3: The number of consumers and those who were not used folic acid

	GROUP		Total
	Control	Case	
Not use b9	8	12	20
Use	47	49	96
Total	55	61	116

In the study of 116 people, 96 subjects received a supplement of folic acid during pregnancy. Forty-nine women were patients with preeclampsia and 47 pregnant women were in the control group. The people who have not been received supplementation during pregnancy were 20. Finally, 60% of them have been diagnosed with gestational hypertension and have been hospitalized, and 40% were in the control group.

DISCUSSION

Pre-eclampsia or gestational hypertension is a multifactorial disorder that increases the sulphur-containing amino acid homocysteine. The aetiology is of general interest, since homocysteine is a marker of vascular disease(40) (41) (42) (43). Some foods contain antioxidants, including vitamins C-E and folic acid, which reduce homocysteine and pre-eclampsia. The increased homocysteine level reduces the amount of antioxidant enzymes, including key elements in maintaining cellular

homeostasis and reducing oxidative damage (44) (45) found in some studies of TAS reduction in women with pre-eclampsia compared with normal pregnant women

(46) The use of proper antioxidants to enhance total antioxidant capacity and reduce vascular disease is recommended (47) Tryptophan-fortified cereals are factors that enhance TAS (48) Studies of the effects of tea and mint, and other herbal extracts, on TAS in women with pre-eclampsia show they reduce oxidative stress and increase TAS in both mother and fetus (49) Many studies on the etiology and severity of preeclampsia have associated with increased levels of homocysteine. Further, a decrease in placental folate and metabolism disorders related to folate have introduced as pathogenic factors in preeclampsia (50) Many studies on the etiology and severity of preeclampsia have associated with increased levels of homocysteine. Further, a decrease in placental folate and metabolism disorders related to folate have introduced as pathogenic factors in preeclampsia (51) Among the factors that are involved in the reduction of folic acid, methylene tetrahydrofolate reductase gene mutation, inadequate intake of food—Sweden absorption and liver disorders can be cited (52) (53) (54) According to the recent studies during preeclampsia homocysteine-induced vascular disease increases, and the role of serum levels of folic acid is significant disagreement (55) Recognising the importance of folic acid in reducing blood pressure much research has been conducted. The study was conducted on mice and the results indicate that the intake of folic acid-rich diet low in methionine increased the ratio of albumin to creatinine and proteinuria, which is one of the symptoms of pre eclampsia (56) Generally, lack of cobalamin and folic acid deficiency increases homocysteine that is a risk factor for vascular disease, including high blood pressure (57) In this regard, a recent study results also suggest that the role of folic acid in reducing the incidence of preeclampsia.

Conclusion

Increased homocysteine leads to oxidative stress, and its effects in women with preeclampsia, and a diet rich in antioxidants, including folic acid supplements reduce the effects of oxidative stress caused by

homocysteine that is considered as a predictive factor for preeclampsia.

AKNOWLEDGEMENT

The authors wish to thank the patients for participating in the study also we are grateful from Drug applied Research Center, Tabriz University of Medical Science for their collaboration.

Conflict of Interest

The authors have declared no conflicts of interest.

References

1. Mahan LK, Escott-Stump S, Raymond JL, Krause MV. Krause's food & the nutrition care process: Elsevier Health Sciences; 2012.
2. Wen SW, Chen X-K, Rodger M, White RR, Yang Q, Smith GN, et al. Folic acid supplementation in early second trimester and the risk of preeclampsia. *American journal of obstetrics and gynecology*. 2008;198(1):45. e1-e7.
3. Polsani S, Phipps E, Jim B. Emerging new biomarkers of preeclampsia. *Advances in chronic kidney disease*. 2013;20(3):271-9.
4. NADAFI M, HOSSEINI M, AFRASIABYFAR A, MOMENI E, MALEKZADEH G. Paper: ASSOCIATION OF HOMOCYSTEINE, VITAMIN AND BLOOD FACTORS WITH PREECLAMPSIA IN PREGNANT WOMEN.
5. Ghulmiyyah L, Sibai B, editors. Maternal mortality from preeclampsia/eclampsia. *Seminars in perinatology*; 2012: Elsevier.
6. Clark SL, Belfort MA, Dildy GA, Herbst MA, Meyers JA, Hankins GD. Maternal death in the 21st century: causes, prevention, and relationship to cesarean delivery. *American journal of obstetrics and gynecology*. 2008;199(1):36. e1-e5.
7. Rahimi G, Tazakori Z, Shateri N. Relation between Homocysteine serum levels and pregnancy complicated with preeclampsia occurrence. *Journal of Ardabil University of Medical Sciences*. 2010;10(2):121-7.
8. Digre KB, Varner MW, Osborn AG, Crawford S. Cranial magnetic resonance imaging in severe preeclampsia vs eclampsia. *Archives of neurology*. 1993;50(4):399-406.
9. Al-Mulhim A-A, Abu-Heija A, Al-Jamma F, El-Harith e-H. Pre-eclampsia: maternal risk factors and perinatal outcome. *Fetal diagnosis and therapy*. 2002;18(4):275-80.

10. Bolander-Gouaille C. Focus on Homocysteine and the Vitamins Involved in its metabolism: Involved in Its Metabolism: Springer Science & Business Media; 2001.
11. Miller AL. The methionine-homocysteine cycle and its effects on cognitive diseases. *Alternative medicine review: a journal of clinical therapeutic*. 2003;8(1):7-19.
12. Rees WD, Wilson FA, Maloney CA. Sulfur amino acid metabolism in pregnancy: the impact of methionine in the maternal diet. *The Journal of nutrition*. 2006;136(6):1701S-5S.
13. Brosnan JT, Brosnan ME. The sulfur-containing amino acids: an overview. *The Journal of nutrition*. 2006;136(6):1636S-40S.
14. Lentz S, Sadler JE. Inhibition of thrombomodulin surface expression and protein C activation by the thrombogenic agent homocysteine. *Journal of clinical investigation*. 1991;88(6):1906.
15. Piccione G, Casella S, Assenza A, Fazio F, Caola G. Evaluation of serum homocysteine and oxidative stress during lactation in ewes. *Czech J Anim Sci*. 2008;53(Suppl 11):462-5.
16. Cavalca V, Cighetti G, Bamonti F, Loaldi A, Bortone L, Novembrino C, et al. Oxidative stress and homocysteine in coronary artery disease. *Clinical chemistry*. 2001;47(5):887-92.
17. Shiraishi M, Haruna M, Matsuzaki M, Ota E, Murayama R, Watanabe E, et al. Association between oxidized LDL and folate during pregnancy. *Biological research for nursing*. 2011;1099800411427581.
18. Amaral LM, Pinheiro LC, Guimaraes DA, Palei AC, Sertório JT, Portella RL, et al. Antihypertensive effects of inducible nitric oxide synthase inhibition in experimental pre-eclampsia. *Journal of cellular and molecular medicine*. 2013;17(10):1300-7.
19. Murphy M. How mitochondria produce reactive oxygen species. *Biochem J*. 2009;417:1-13.
20. Halliwell B, Gutteridge J. Role of free radicals and catalytic metal ions in human disease: an overview. *Methods in enzymology*. 1990;186:1.
21. Halliwell B. Free radicals and other reactive species in disease: Wiley Online Library; 2005.
22. Murphy MP. Mitochondrial dysfunction indirectly elevates ROS production by the endoplasmic reticulum. *Cell metabolism*. 2013;18(2):145-6.
23. Kukat A, Dogan SA, Edgar D, Mourier A, Jacoby C, Maiti P, et al. Loss of ucp2 attenuates mitochondrial dysfunction without altering ros production and uncoupling activity. *PLoS genetics*. 2014;10(6):e1004385.
24. Yamamoto YY. Free radicals in chemistry, biology and medicine: OICA International (UK); 2000.
25. Rogers EJ, Chen S, Chan A. Folate deficiency and plasma homocysteine during increased oxidative stress. *New England Journal of Medicine*. 2007;357(4):421-2.
26. Jordao Júnior AA, Domenici FA, Lатарo RC, Portari GV, Vannucchi H. Effect of methionine load on homocysteine levels, lipid peroxidation and DNA damage in rats receiving ethanol. *Brazilian Journal of Pharmaceutical Sciences*. 2009;45(4):709-14.
27. Kashinakunti S, Sunitha H, Gurupadappa K, Shankarprasad D, Suryaprakash G, Ingin J. Lipid peroxidation and antioxidant status in preeclampsia. *Al Ameen J Med Sci*. 2010;3(1):38-41.
28. Mujawar SA, Patil VW, Daver RG. Study of serum homocysteine, folic acid and vitamin B12 in patients with preeclampsia. *Indian journal of clinical biochemistry*. 2011;26(3):257-60.
29. Villar J, Merialdi M, Gülmezoglu AM, Abalos E, Carroli G, Kulier R, et al. Nutritional interventions during pregnancy for the prevention or treatment of maternal morbidity and preterm delivery: an overview of randomized controlled trials. *The Journal of nutrition*. 2003;133(5):1606S-25S.
30. Januszeski S. When Roots Intertwine: The Language and Politics of Raw Foodism: Citeseer; 2013.
31. Nilsson K, Gustafson L, Hultberg B, et al. Plasma homocysteine is a sensitive marker for tissue deficiency of both cobalamines and folates in a psychogeriatric population. *Dementia and geriatric cognitive disorders*. 1998;10(6):476-82.
32. Scholl TO, Johnson WG. Folic acid: influence on the outcome of pregnancy. *The American journal of clinical nutrition*. 2000;71(5):1295s-303s.
33. Selhub J, Miller JW. The pathogenesis of homocysteinemia: interruption of the coordinate regulation by S-adenosylmethionine of the remethylation and transsulfuration of homocysteine. *The American journal of clinical nutrition*. 1992;55(1):131-8.
34. Ueland PM, Refsum H. Plasma homocysteine, a risk factor for vascular disease: plasma levels in health, disease, and drug therapy. *J Lab Clin Med*. 1989;114(5):473-501.
35. Shils ME, Shike M. Modern nutrition in health and disease: Lippincott Williams & Wilkins; 2006.
36. Cattaneo M. Hyperhomocysteinemia, atherosclerosis and thrombosis. *Thromb Haemost*. 1999;81(2):165-76.

37. Abraham JM, Cho L. The homocysteine hypothesis: still relevant to the prevention and treatment of cardiovascular disease? *Cleveland Clinic journal of medicine*. 2010;77(12):911-8.
38. Balk EM, Raman G, Tatsioni A, Chung M, Lau J, Rosenberg IH. Vitamin B6, B12, and folic acid supplementation and cognitive function: a systematic review of randomized trials. *Archives of Internal Medicine*. 2007;167(1):21-30.
39. Willett WC, Stampfer MJ. What vitamins should I be taking, doctor? *New England Journal of Medicine*. 2001;345(25):1819-24.
40. Rumbold A, Duley L, Crowther CA, Haslam RR. Antioxidants for preventing pre-eclampsia. *The Cochrane Library*. 2008.
41. Polyzos NP, Mauri D, Tsappi M, Tzioras S, Kamosioras K, Cortinovis I, et al. Combined vitamin C and E supplementation during pregnancy for preeclampsia prevention: a systematic review. *Obstetrical & gynecological survey*. 2007;62(3):202-6.
42. Rumbold A, Crowther CA. Vitamin E supplementation in pregnancy. *The Cochrane Library*. 2005.
43. Conde-Agudelo A, Romero R, Kusanovic JP, Hassan SS. Supplementation with vitamins C and E during pregnancy for the prevention of preeclampsia and other adverse maternal and perinatal outcomes: a systematic review and metaanalysis. *American journal of obstetrics and gynecology*. 2011;204(6):503.e1-. e12.
44. Mani M, Golmohammadi T, Khaghani S, Zamani Z, Azadmanesh K, Meshkani R, et al. Homocysteine Induces Heme Oxygenase-1 Expression via Transcription Factor Nrf2 Activation in HepG2 Cells. *Iranian biomedical journal*. 2013;17(2):93.
45. Brito JA, do Nascimento Marreiro D, Neto JMM, e Silva DMC, de Sousa Almondes KG, Neto JdDV, et al. Enzyme activity of superoxide dismutase and zincemia in women with preeclampsia. *Nutr Hosp*. 2013;28(2):486-90.
46. Gupta S, Aziz N, Sekhon L, Agarwal R, Mansour G, Li J, et al. Lipid peroxidation and antioxidant status in preeclampsia: a systematic review. *Obstetrical & gynecological survey*. 2009;64(11):750-9.
47. Bravo R, Matito S, Cubero J, Paredes S, Franco L, Rivero M, et al. Tryptophan-enriched cereal intake improves nocturnal sleep, melatonin, serotonin, and total antioxidant capacity levels and mood in elderly humans. *Age*. 2013;35(4):1277-85.
48. Padmini E, Usha Rani M, Lavanya S. Effect of mint and tea infusions on the antioxidant capacity of preeclampsic endothelial cells. *Asian Journal of Microbiology, Biotechnology and Environmental Science*. 2008;10(4):903-9.
49. Ray J, Diamond P, Singh G, Bell C. Brief overview of maternal triglycerides as a risk factor for pre- eclampsia. *BJOG: An International Journal of Obstetrics & Gynaecology*. 2006;113(4):379-86.
50. Mislanova C, Martsenyuk O, Huppertz B, Obolenskaya M. Placental markers of folate-related metabolism in preeclampsia. *Reproduction*. 2011;142(3):467-76.
51. Giugliani E, Jorge S, Goncalves A. Serum and red blood cell folate levels in parturients, in the intervillous space of the placenta and in full-term newborns. *Journal of Perinatal Medicine-Official Journal of the WAPM*. 1985;13(2):55-60.
52. Giugliani ER, Jorge S, Goncalves A. Serum vitamin B12 levels in parturients, in the intervillous space of the placenta and in full-term newborns and their interrelationships with folate levels. *The American journal of clinical nutrition*. 1985;41(2):330-5.
53. Cetin I. Amino acid interconversions in the fetal-placental unit: the animal model and human studies in vivo. *Pediatric research*. 2001;49(2):148-54.
54. Camelo Jr J, Jorge S, Martinez F. Amino acid composition of parturient plasma, the intervillous space of the placenta and the umbilical vein of term newborn infants. *Brazilian journal of medical and biological research*. 2004;37(5):711-7.
55. Daly S, Cotter A, Molloy AE, Scott J, editors. Homocysteine and folic acid: implications for pregnancy. *Seminars in vascular medicine*; 2005.
56. Falcao S, Bisotto S, Gutkowska J, Lavoie JL. Hyperhomocysteinemia is not sufficient to cause preeclampsia in an animal model: the importance of folate intake. *American journal of obstetrics and gynecology*. 2009;200(2):198. e1-. e5.
57. Klee GG. Cobalamin and folate evaluation: measurement of methylmalonic acid and homocysteine vs vitamin B12 and folate. *Clinical chemistry*. 2000;46(8):1277-83.