

The Effect of Yoga on Stress, Anxiety, and Depression in Women

Abstract

Background: In recent decades, several medical and scientific studies on yoga proved it to be very useful in the treatment of some diseases. This study was conducted to investigate the effects of yoga on stress, anxiety, and depression in women living in Ilam, Iran. **Methods:** This study is a quasi-experimental study with pre-post test. To collect data, the questionnaire of DASS-21 (Depression Anxiety Stress Scale-21) was used. For eligible samples, hatha yoga exercises and training sessions were held for 4 weeks (3 time/weeks; 60-70 min each) by a specialist. Data were analyzed using SPSS version 20. **Results:** 52 women with a mean age of 33.5 ± 6.5 were included for analysis. Depression, anxiety, and stress decreased significantly in women after 12 sessions of regular hatha yoga practice ($P < 0.001$). **Conclusions:** Yoga has an effective role in reducing stress, anxiety, and depression. Thus, it can be used as complementary medicine.

Keywords: Anxiety, depression, stress, women, yoga

Introduction

Complementary medicine refers to a category of treatments and interventions that have not been raised in modern medicine.^[1] Yoga is a Sanskrit word meaning unity of mind and body, which has been used in Eastern societies since 5000 years ago and has recently received much attention from Western countries.^[2] In recent decades, several medical and scientific studies on yoga proved it to be very useful in the treatment of some diseases.^[3] Studies have demonstrated effect of yoga for many conditions, including multiple sclerosis,^[4] asthma,^[5] irritable bowel syndrome,^[6] lymphoma,^[7] hypertension,^[8] drug addiction,^[9] osteoarthritis,^[10] and mental health issues.^[11]

Increased stress, depression and anxiety are the features of modern lifestyle.^[12] Due to the adverse effects of drugs in the treatment of anxiety and depression, and in some cases their lack of effectiveness, researchers seek nonpharmacological and noninvasive treatment for these disorders.^[13] Yoga exercises was improved the variables of self-description, psychological status, and the quality of life.^[14] Researches suggest that yoga as an intellectual and mental exercise, improves health feeling.^[15] Furthermore, yoga can improve

the psychological conditions for monitoring and managing stress and negative emotions, increase positive emotions, and help mental balance.^[11] However, despite the popularity and the positive psychological and physiological effects of yoga, it is not widely studied to find how much it really prevents and treats mental disorders. Research in this area is limited in Iran. This study was conducted to investigate the effects of yoga on stress, anxiety, and depression in women living in Ilam from Iran, during 2014 to 2015.

Methods

Design and ethics

This study is a quasi-experimental study with pre-post test that was approved by the Ethics Committee of the Ilam University of Medical Sciences. The population consisted of all women admitted to yoga club in the city of Ilam in 2014–2015.

Inclusion and exclusion criteria

Inclusion criteria were educated, nonathlete and nonpregnant women with the ability to perform hatha yoga exercises without inability to exercise. Exclusion criteria included refusal or unwillingness to perform yoga continuously, simultaneous exercise, and receiving medication for mental disorders.

How to cite this article: Shohani M, Badfar G, Nasirkandy MP, Kaikhavani S, Rahmati S, Modmeli Y, et al. The effect of yoga on stress, anxiety and depression in women. *Int J Prev Med* 2018;9:21.

Masoumeh Shohani,
Gholamreza
Badfar¹,
Marzieh Parizad
Nasirkandy²,
Sattar Kaikhavani³,
Shoboo Rahmati⁴,
Yaghoob Modmeli⁵,
Ali Soleymani⁶,
Milad Azami⁷

Department of Nursing, Faculty of Allied Medical Sciences, Ilam University of Medical Sciences, Ilam, Iran, ¹Department of Pediatrics, Behbahan School of Medicine, Ahvaz Jundishapur University of Medical Science, Behbahan, Iran, ²Department of Obstetrics and Gynecology, Women's Reproductive Health Research Center, School of Medicine, Tabriz University of Medical Sciences, Tabriz, Iran, ³Prevention of Psychosocial Injuries Research Center, Faculty of medicine, Ilam University of Medical Sciences, Ilam, Iran, ⁴M.Sc. Student of Epidemiology, Student Research Committee, School of Health, Ilam University of Medical Sciences, Ilam, Iran, ⁵M.Sc. Student of Nursing, Student Research Committee, Ahvaz Jundishapur University of Medical Sciences, Ahvaz, Iran, ⁶M.Sc. in Management, Department of Finance, School of Medicine, Dezfoul University of Medical Sciences, Dezfoul, Iran, ⁷Medical Student, Student Research Committee, Ilam University of Medical Sciences, Ilam, Iran

Address for correspondence:

Dr. Milad Azami,
Student Research Committee,
Ilam University of Medical
Sciences, Ilam, Iran.
E-mail: MiladAzami@medilam.ac.ir

Access this article online

Website:
www.ijpvmjournal.net/www.ijpvm.ir

DOI:
10.4103/ijpvm.IJPVM_242_16

Quick Response Code:



This is an open access article distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as the author is credited and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Assessment tools and management

To collect data, the questionnaire of DASS-21 (Depression Anxiety Stress Scale-21) was used. The validity and reliability of this standard questionnaire was examined by Sahebi *et al.* and Cronbach's alpha was estimated 0.7, 0.66 and 0.76 for depression, anxiety, and stress, respectively. in a study entitled "validation of depression anxiety and stress scale for an Iranian population".^[16] Each of the above mentioned states are assessed with seven questions. Hatha yoga exercises and training sessions were held 3 time/weeks; 60-70 min each (postures, breathing techniques, meditation) by a specialist. Before the intervention, questionnaires were completed by women. The intervention lasted 12 sessions. At the end of the 12th session, the questionnaire of DASS-21 was again completed by women.

Statistical procedures

The obtained data were analyzed using SPSS version 20 (IBM, Armonk, NY, USA). According to the established normality, paired sample *t*-test was used for comparing the results before and after the intervention. The threshold of significance was set at $P < 0.05$.

Results

The total eligible sample consisted of 52 women with a mean age of 33.5 ± 6.5 years. Other demographic characteristics are shown in Table 1.

The difference between mean scores of depression, anxiety, and stress before and after 12 sessions of regular hatha yoga practice was statistically significant [Table 2].

Discussion

The present study showed that 12 sessions of intervention as regular hatha yoga exercise significantly reduced stress, anxiety, and depression in women.

In other studies, including Tayyebi *et al.*, among hemodialysis patients,^[17] Rahnama *et al.*, among multiple sclerosis patients,^[15] Javnbakht *et al.*, among women living in Mashhad,^[18] and Gong among pregnant women,^[19] the significant positive effects of yoga in reducing stress, anxiety, and depression have been confirmed. In the study of Streeter *et al.*,^[20] it was found that greater improvement in mood and greater decreases in anxiety during 12 week of yoga intervention compared to walking group. Oken *et al.*^[4] did not observe any significant effect of yoga on improvement in mood in patients with MS, the limitation of this study is the small number of intervention sessions (one session per week).

Dalgas *et al.* proved that the nature of yoga is controlling the mind and central nervous system and unlike other sports, it has a moderating effect on the nervous system, the hormonal emissions, physiological factors, and regulation of nerve impulses; therefore, it can be effective in improving depression and mental disorders.^[21]

Table 1: Demographic characteristics of the women

Variable	n (%)
Age	33.5 ± 6.5*
Marital status (%)	
Single	18 (34.6)
Married	34 (65.4)
Job (%)	
Unemployed	10 (19.2)
Housewife	19 (36.5)
Employed	9 (17.3)
Retired	2 (3.8)
Other jobs	12 (23.1)
Level of income (%)	
Low	6 (11.5)
Average	36 (69.2)
High	10 (19.2)
Education (%)	
Primary	3 (5.8)
Diploma	13 (25)
University degrees	36 (69.2)

*Mean±SD. SD=Standard deviation

Table 2: A comparison of the mean scores of stress, anxiety, and depression before and after the intervention of 12 sessions of regular hatha yoga practice session (n=52)

Variable	n	Mean±SD	Correlation	P
Depression				
Before intervention	52	6.5 (5.5)	0.8	0.001
After intervention	52	5.1 (5)		
Anxiety				
Before intervention	52	5.7 (4.5)	0.7	0.0001
After intervention	52	4.2 (4)		
Stress				
Before intervention	52	7.8 (4.8)	0.7	0.0001
After intervention	52	5.6 (4.2)		

SD=Standard deviation

One limitation of this study was that only the women were studied and since there were no yoga clubs for men, they have not been studied in this research.

Conclusions

Yoga has an effective role in reducing stress, anxiety, and depression that can be considered as complementary medicine and reduce the medical cost per treatment by reducing the use of drugs. Given that, the reason behind the effect of yoga on stress, anxiety, and depression is not clear for us and may be transient, and it is suggested that future studies are done to investigate the long-term effect of yoga on stress, anxiety, and depression.

Acknowledgments

The authors would like to thank Ilam University of Medical Sciences for financial support and also yoga instructors (Miss. Morshedi and Miss. Khosravi) for their cooperation.

Financial support and sponsorship

Ilam University of Medical Sciences.

Conflicts of interest

There are no conflicts of interest.

Received: 24 Jul 16 **Accepted:** 14 Jun 17

Published: 21 Feb 18

References

1. Borji M, Otaghi M, Salimi E, Sanei P. Investigating the effect of performing the quiet time protocol on the sleep quality of cardiac patients. *Biomedical Research* 2017;28:7076-80.
2. Barnes PM, Powell-Griner E, McFann K, Nahin RL. Complementary and alternative medicine use among adults: United States, 2002. *Adv Data* 2004;1-19.
3. Singh S, Malhotra V, Singh KP, Sharma SB, Madhu SV, Tandon OP. A preliminary report on the role of yoga asanas on oxidative stress in non-insulin dependent diabetes mellitus. *Indian J Clin Biochem* 2001;16:216-20.
4. Oken BS, Kishiyama S, Zajdel D, Bourdette D, Carlsen J, Haas M, *et al.* Randomized controlled trial of yoga and exercise in multiple sclerosis. *Neurology* 2004;62:2058-64.
5. Freitas DA, Holloway EA, Bruno SS, Chaves GS, Fregonezi GA, Mendonça KP. Breathing exercises for adults with asthma. *Cochrane Database Syst Rev* 2013;1:CD001277.
6. Taneja I, Deepak KK, Poojary G, Acharya IN, Pandey RM, Sharma MP. Yogic versus conventional treatment in diarrhea-predominant irritable bowel syndrome: A randomized control study. *Appl Psychophysiol Biofeedback* 2004;29:19-33.
7. Cohen L, Warneke C, Fouladi RT, Rodriguez MA, Chaoul-Reich A. Psychological adjustment and sleep quality in a randomized trial of the effects of a Tibetan yoga intervention in patients with lymphoma. *Cancer* 2004;100:2253-60.
8. Chu P, Gotink RA, Yeh GY, Goldie SJ, Hunink MG. The effectiveness of yoga in modifying risk factors for cardiovascular disease and metabolic syndrome: A systematic review and meta-analysis of randomized controlled trials. *Eur J Prev Cardiol* 2016;23:291-307.
9. Shaffer HJ, LaSalvia TA, Stein JP. Comparing Hatha yoga with dynamic group psychotherapy for enhancing methadone maintenance treatment: A randomized clinical trial. *Altern Ther Health Med* 1997;3:57-66.
10. Garfinkel MS, Schumacher HR Jr, Husain A, Levy M, Reshetar RA. Evaluation of a yoga based regimen for treatment of osteoarthritis of the hands. *J Rheumatol* 1994;21:2341-3.
11. Duan-Porter W, Coeytaux RR, McDuffie JR, Goode AP, Sharma P, Mennella H, *et al.* Evidence map of yoga for depression, anxiety, and posttraumatic stress disorder. *J Phys Act Health* 2016;13:281-8.
12. Brandon H Hidaka, B.A. Depression as a disease of modernity: explanations for increasing prevalence. *J Affect Disord*. 2012;140:205-14.
13. Little N. Depression Treatment Options. Available from: <http://www.insightjournal.com/>. [Last cited on 2007 Jan 01].
14. Richter S, Tietjens M, Ziereis S, Querfurth S, Jansen P. Yoga Training in Junior Primary School-Aged Children Has an Impact on Physical Self-Perceptions and Problem-Related Behavior. *Frontiers in Psychology* 2016;7:203.
15. Rahnama N, Bambaiechi E, Arbabzadeh S, Sadeghipour H, Etemadifar M, Namazizadeh M. Effects of yoga on depression in women with multiple sclerosis. *J Isfahan Med Sch* 2011;29:483-90.
16. Sahebi A, Asghari A, Salari RS. [Validation of depression anxiety and stress scale for an Iranian population] *Journal of Iranian psychologists* 2005;1:299-312. (Persian).
17. Tayyebi A, Babahaji M, Sadeghishermeh M, Ebadi A, Eynollahi B. Study of the effect of Hatha Yoga exercises on stress, anxiety and depression among hemodialysis patients. *IJCCN* 2011;4:67-72.
18. Javnbakht M, Hejazi Kenari R, Ghasemi M. Effects of yoga on depression and anxiety of women. *Complement Ther Clin Pract* 2009;15:102-4.
19. Gong H, Ni C, Shen X, Wu T, Jiang C. Yoga for prenatal depression: A systematic review and meta-analysis. *BMC Psychiatry* 2015;15:14.
20. Streeter CC, Whitfield TH, Owen L, Rein T, Karri SK, Yakhkind A, *et al.* Effects of yoga versus walking on mood, anxiety, and brain GABA levels: A randomized controlled MRS study. *J Altern Complement Med* 2010;16:1145-52.
21. Dalgas U, Stenager E, Ingemann-Hansen T. Multiple sclerosis and physical exercise: Recommendations for the application of resistance-, endurance- and combined training. *Mult Scler* 2008;14:35-53.